

HENRY

Hydraulic Engineering Repository

Ein Service der Bundesanstalt für Wasserbau

Conference Paper, Published Version

Verschoren, Veerle; Schwarz, Christian; Schoelynck, Jonas; Buis, Kerst; Antoine, Germain; Claude, Nicolas; Meire, Patrick; Temmerman, Stijn
Implementing plant growth of flexible aquatic vegetation into a hydrodynamic model (TELEMAC2D)

Zur Verfügung gestellt in Kooperation mit/Provided in Cooperation with:
TELEMAC-MASCARET Core Group

Verfügbar unter/Available at: <https://hdl.handle.net/20.500.11970/104337>

Vorgeschlagene Zitierweise/Suggested citation:

Verschoren, Veerle; Schwarz, Christian; Schoelynck, Jonas; Buis, Kerst; Antoine, Germain; Claude, Nicolas; Meire, Patrick; Temmerman, Stijn (2015): Implementing plant growth of flexible aquatic vegetation into a hydrodynamic model (TELEMAC2D). In: Moulinec, Charles; Emerson, David (Hg.): Proceedings of the XXII TELEMAC-MASCARET Technical User Conference October 15-16, 2036. Warrington: STFC Daresbury Laboratory. S. 213-214.

Standardnutzungsbedingungen/Terms of Use:

Die Dokumente in HENRY stehen unter der Creative Commons Lizenz CC BY 4.0, sofern keine abweichenden Nutzungsbedingungen getroffen wurden. Damit ist sowohl die kommerzielle Nutzung als auch das Teilen, die Weiterbearbeitung und Speicherung erlaubt. Das Verwenden und das Bearbeiten stehen unter der Bedingung der Namensnennung. Im Einzelfall kann eine restriktivere Lizenz gelten; dann gelten abweichend von den obigen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Documents in HENRY are made available under the Creative Commons License CC BY 4.0, if no other license is applicable. Under CC BY 4.0 commercial use and sharing, remixing, transforming, and building upon the material of the work is permitted. In some cases a different, more restrictive license may apply; if applicable the terms of the restrictive license will be binding.



Implementing plant growth of flexible aquatic vegetation into a hydrodynamic model (TELEMAC2D)

Veerle Verschoren¹, Christian Schwarz¹, Jonas Schoelynck¹, Kerst Buis¹, Germain Antoine^{2,3}, Nicolas Claude^{2,3}, Patrick Meire¹, Stijn Temmerman¹

¹ University of Antwerp, Universiteitsplein 1, 2610 Antwerp, Belgium

² EDF R&D, National Laboratory for Hydraulics and Environment (LNHE) & Saint Venant Laboratory for Hydraulics, Chatou, France

³ Saint-Venant Hydraulics Laboratory, Université Paris-Est, Chatou, France
Corresponding author: Veerle Verschoren, veerle.verschoren@uantwerpen.be

Proposed session: Water quality, ecology and environmental impact

Keywords: influence of aquatic vegetation, coupling of code, hydraulic resistance

Speaker: Veerle Verschoren

Abstract:

In lowland rivers, mutual interactions between aquatic vegetation and water flow are expected to impact morphology. The presence of aquatic vegetation increases the hydraulic resistance encountered by water flow. On a local scale this results in flow deceleration within vegetation patches and in flow acceleration adjacent to vegetation patches. Reduced flow velocities within vegetation patches are beneficial for plant growth and therefore referred to as a positive feedback. While increased velocities adjacent to the vegetation patches are able to hamper plant growth, break off or uproot plants constituting a negative feedback. These two phenomena are in literature referred to as scale-dependent feedbacks (positive and negative feedbacks dependent on the spatial scale considered, i.e. in the patch or at its edges), which are further described as the mechanisms influencing heterogeneous spatial habitat structure.

A depth-averaged hydrodynamic model (TELEMAC2D) coupled with a plant model is used to investigate the importance of scale-dependent feedbacks in structuring lowland river habitats. We simulate the feedback between spatio-temporal growth of aquatic vegetation and its impact on the associated flow field. The dynamic plant growth model simulates changes in biomass of the plants. The biomass is represented as a tracer in the hydrodynamic model. The plant model itself consists of vegetation establishment-, mortality-, logistic increase of biomass- and lateral spatial expansion processes. The influence of vegetation is incorporated by schematizing vegetation as cylinders which exert a drag force to the flow, following the approach of Baptist et al. (2007). Since vegetation is flexible, the magnitude of the drag force is a function of vegetation reconfiguration and therefore dependent on the stream velocity itself.

Our preliminary results stress the importance of scale-dependent feedbacks on spatial habitat structure. We focus on three species with contrasting plant traits which due to differences in scale-dependent feedback strength lead to different spatial configurations. A dense species (*Callitriche obtusangula*) causing high hydraulic resistance instigating a strong scale-dependent feedback which results in delineated patches alternated with bare soil. An intermediate dense species (*Sparganium emersum*) forms long vegetated areas and a low density species (*Potamogeton natans*) with low hydraulic resistance, limited interaction with the water flow occupies the complete river bed. Our results underline, through a comparison between our model results and observed vegetation

patterns in lowland rivers, the importance of species-specific plant traits on spatial habitat structure.

References:

Baptist, M., V. Babovic, J. Rodríguez Uthurburu, M. Keijzer, R. Uittenbogaard, A. Mynett, and A. Verwey (2007), On inducing equations for vegetation resistance, *Journal of Hydraulic Research*, 45(4), 435-450.